
Function Of The Cardiac Sphincter

*Function Of The
Cardiac Sphincter*

Downloaded from turn-nyc-edge-vdmdp.louper.io by
Ximena & Morrow

UNDERSTANDING THE CARDIAC SPHINCTER: A VITAL GUARDIAN OF DIGESTIVE HEALTH

Deep within the human body, a small but mighty muscular ring performs a critical role every single moment of the day. The cardiac sphincter, also widely known as the lower esophageal sphincter (LES), acts as a sophisticated gatekeeper between the esophagus and the stomach. Despite its modest size, this

anatomical structure is responsible for one of the most important functions in digestion: ensuring that food moves in the right direction. Understanding the function of the cardiac sphincter is essential for anyone interested in digestive health, acid reflux prevention, and overall well-being.

The term cardiac sphincter might sound like it has something to do with the heart, and indeed it does in a roundabout way. The name derives from the Greek word kardia, meaning heart, because this sphincter is located near

the cardia of the stomach, which lies close to the heart. However, its function is entirely digestive. In modern medical terminology, lower esophageal sphincter is the preferred clinical term, but cardiac sphincter remains widely used in anatomy and physiology contexts. This article will explore every aspect of this essential structure, from its anatomy and primary roles to the consequences when it malfunctions.

ANATOMY AND LOCATION OF THE CARDIAC SPHINCTER

The cardiac sphincter is located at the junction where the esophagus meets the stomach. More precisely, it sits at the gastroesophageal junction, approximately 40 centimeters from the upper incisor teeth in an average adult.

This region is anatomically complex because it represents a transition point between two very different environments: the neutral pH of the esophagus and the highly acidic environment of the stomach.

The sphincter itself is not a distinct, separate anatomical structure like a valve in a pipe. Rather, it is a zone of specialized circular smooth muscle fibers that are tonically contracted at rest. This region is approximately 2 to 4 centimeters in length and is reinforced by several additional anatomical features, including the diaphragmatic crura and the phrenoesophageal ligament. These surrounding structures work in concert with the smooth muscle to maintain closure pressure. The intrinsic smooth muscle of the cardiac

sphincter is unique because it maintains a state of partial contraction even when at rest, a property known as resting tone. This constant tension is what prevents stomach contents from refluxing into the esophagus.

THE PRIMARY FUNCTION OF THE CARDIAC SPHINCTER

Preventing Gastroesophage al Reflux

The single most important function of the cardiac sphincter is to prevent the backward flow of stomach contents into

the esophagus. This is a remarkable task because the stomach is a turbulent environment. It churns food, secretes hydrochloric acid with a pH as low as 1.5, and generates significant pressure during digestion. The esophagus, by contrast, has delicate mucosal tissue that cannot tolerate exposure to gastric acid. By maintaining a high-pressure zone at the gastroesophageal junction, the cardiac sphincter acts as a barrier that keeps acid, bile, partially digested food, and digestive enzymes where they belong, inside the stomach. Without this barrier, reflux would occur constantly, leading to severe esophagitis and damage to the esophageal lining.

Regulating Food Passage Into the Stomach

Beyond its barrier function, the cardiac sphincter also plays an active role in allowing food and liquids to pass from the esophagus into the stomach. When a person swallows, a complex sequence of events unfolds. The upper esophageal sphincter relaxes to accept the bolus, and peristaltic waves propel the food downward. As the bolus approaches the lower esophagus, the cardiac sphincter receives signals via the vagus nerve to relax. This relaxation is mediated by inhibitory neurotransmitters, primarily

nitric oxide and vasoactive intestinal peptide. The sphincter opens widely enough to permit passage of the food bolus, then promptly closes again to reestablish the barrier. This process is known as deglutitive relaxation and typically lasts only 5 to 10 seconds. The precise timing and coordination of this relaxation are essential. If the sphincter fails to relax properly, a condition called esophageal dysphagia results, where food feels stuck in the chest or throat.

HOW THE CARDIAC SPHINCTER FUNCTIONS IN DETAIL

Resting Tone and

Pressure Dynamics

At rest, the cardiac sphincter maintains a pressure of approximately 15 to 30 mmHg above gastric pressure. This pressure gradient is what keeps the barrier intact. The resting tone is generated by the smooth muscle itself and is influenced by hormonal, neural, and mechanical factors. For example, gastrin and motilin are hormones that increase sphincter pressure, while secretin and cholecystokinin decrease it. The vagus nerve provides both excitatory and inhibitory input. Sympathetic activation, such as during stress, can also modulate sphincter tone.

This intricate regulation allows the body to adjust barrier strength based on digestive demands and overall physiological state.

Transient Lower Esophageal Sphincter Relaxations

One of the most fascinating aspects of cardiac sphincter function is the phenomenon known as transient lower esophageal sphincter relaxation (TLESR). These are brief, spontaneous relaxations of the sphincter that are not triggered by

swallowing. TLESRs are the primary mechanism by which gas is vented from the stomach, allowing belching to occur. However, they are also the main cause of acid reflux episodes, especially in individuals with gastroesophageal reflux disease (GERD). In healthy individuals, TLESRs occur several times per hour, but the volume of refluxed material is minimal and rapidly cleared by esophageal peristalsis and saliva. In individuals with GERD, TLESRs are more frequent and often associated with larger volumes of acid reflux, leading to symptoms and tissue damage.

The Role of the

Diaphragm in Sphincter Function

The diaphragmatic crura, muscle fibers from the diaphragm that surround the esophagus, provide additional external compression to the cardiac sphincter. This is particularly important during inspiration, when negative thoracic pressure and positive abdominal pressure would otherwise promote reflux. The crural diaphragm contracts during inspiration, increasing pressure on the lower esophagus and reinforcing the barrier. This coordination between the smooth muscle of the sphincter and

the skeletal muscle of the diaphragm is a brilliant example of how different muscle types work together. During a TLESR, the crural diaphragm also relaxes, allowing gas to escape. When this coordination is disrupted, such as in a hiatal hernia, the barrier becomes less effective and reflux becomes more likely.

FACTORS THAT AFFECT CARDIAC SPHINCTER FUNCTION

Numerous lifestyle, dietary, and medical factors can influence the function of the cardiac sphincter. Understanding these factors is key to managing or preventing dysfunction.

- **Dietary factors:** High-fat meals, chocolate, peppermint, caffeine,

alcohol, and spicy foods can decrease lower esophageal sphincter pressure. Fatty foods, in particular, delay gastric emptying and stimulate the release of cholecystokinin, which relaxes the sphincter.

- **Obesity:** Excess abdominal fat increases intra-abdominal pressure, which overwhelms the cardiac sphincter and promotes reflux. This is one of the strongest risk factors for GERD.
- **Pregnancy:** Hormonal changes, especially increased progesterone, relax smooth muscle throughout the body, including the cardiac sphincter. Combined with increased abdominal pressure from the growing uterus, reflux is

extremely common during pregnancy.

- **Medications:** Several drugs can reduce sphincter pressure, including calcium channel blockers, nitrates, theophylline, benzodiazepines, and anticholinergics. Patients taking these medications for other conditions may develop reflux as a side effect.
- **Smoking:** Nicotine directly relaxes the lower esophageal sphincter and also reduces saliva production, which helps clear acid from the esophagus. Smoking is a well-established risk factor for GERD.
- **Hiatal hernia:** This anatomical abnormality occurs when part of

the stomach pushes upward through the diaphragmatic hiatus. It disrupts the normal positioning and function of the cardiac sphincter, significantly increasing the risk of reflux.

CONDITIONS RELATED TO CARDIAC SPHINCTER DYSFUNCTION

Gastroesophageal Reflux Disease

The most common condition associated with cardiac sphincter dysfunction is GERD. When the sphincter becomes weak, relaxes too frequently, or is

anatomically compromised, gastric acid and contents reflux into the esophagus. Symptoms include heartburn, regurgitation, chest pain, dysphagia, and chronic cough. Over time, chronic GERD can lead to complications such as esophagitis, esophageal strictures, Barrett's esophagus, and an increased risk of esophageal adenocarcinoma. The function of the cardiac sphincter is central to the pathophysiology of GERD, and most treatment strategies aim to either reduce acid secretion or improve barrier function.

Esophageal

Achalasia

Achalasia is a rare disorder in which the cardiac sphincter fails to relax properly during swallowing due to loss of inhibitory neurons in the esophageal myenteric plexus. Patients experience progressive dysphagia, regurgitation of undigested food, chest pain, and weight loss. The condition is essentially the opposite of GERD, where the sphincter is too tight rather than too loose. Diagnosis is confirmed with esophageal manometry, which shows elevated lower esophageal sphincter pressure and absent peristalsis. Treatment options include pneumatic dilation, laparoscopic Heller myotomy, and peroral endoscopic myotomy, all of which aim to disrupt the

non-relaxing sphincter.

Hiatal Hernia

A hiatal hernia occurs when a portion of the stomach protrudes through the diaphragmatic hiatus into the chest cavity. This anatomical displacement separates the intrinsic lower esophageal sphincter from the crural diaphragm, disrupting their coordinated function. There are two main types: sliding hiatal hernia, where the gastroesophageal junction moves above the diaphragm, and paraesophageal hernia, where part of the stomach herniates alongside the esophagus. Both types can impair cardiac sphincter function and lead to chronic reflux. Hiatal hernia is very common, especially in older adults and

those with obesity.

DIAGNOSIS AND ASSESSMENT OF CARDIAC SPHINCTER FUNCTION

When dysfunction is suspected, several diagnostic tools are available to evaluate the cardiac sphincter.

1. **Esophageal manometry:** This is the gold standard for assessing lower esophageal sphincter pressure, relaxation, and coordination. A thin catheter with pressure sensors is passed through the nose into the esophagus, and the patient performs swallows while pressures are recorded. High-resolution manometry provides detailed topographic maps of esophageal

function.

2. **pH monitoring:** To quantify acid reflux, a pH probe is placed in the esophagus, typically 5 cm above the lower esophageal sphincter. The patient wears the probe for 24 hours while recording symptoms and activities. This test correlates reflux episodes with symptoms and provides a DeMeester score.
3. **Impedance monitoring:** Often combined with pH monitoring, impedance detects both acid and non-acid reflux episodes by measuring changes in electrical conductivity between electrodes along the catheter.
4. **Upper endoscopy:** Direct visualization of the gastroesophageal junction allows

assessment of anatomical abnormalities, esophagitis, Barrett's esophagus, and hiatal hernia.

5. **Barium swallow:** This radiographic study shows the anatomy and function of the esophagus and stomach during swallowing, helping to identify strictures, achalasia, and hiatal hernia.

STRATEGIES FOR MAINTAINING A HEALTHY CARDIAC SPHINCTER

While some factors affecting sphincter function are beyond individual control, many lifestyle modifications can support optimal function and reduce the risk of GERD.

- **Maintain a healthy weight:** Reducing excess abdominal fat decreases intra-abdominal pressure and takes strain off the cardiac sphincter. Even modest weight loss can significantly improve reflux symptoms.
- **Eat smaller, more frequent meals:** Large meals distend the stomach and increase gastric pressure, which can overcome sphincter resistance. Eating smaller portions reduces pressure on the barrier.
- **Avoid trigger foods:** Common triggers include fatty foods,

chocolate, coffee, alcohol, peppermint, spicy foods, and citrus. Individual responses vary, so keeping a symptom diary can help identify personal triggers.

- **Elevate the head of the bed:** When lying down, gravity no longer helps keep stomach contents down. Elevating the head of the bed by 15 to 20 centimeters reduces nighttime reflux significantly.
- **Avoid eating before bedtime:** Ideally, finish meals at least three hours before lying down to allow for