
Important Formulas For Algebra 2

LINEAR FUNCTIONS AND EQUATIONS Downloaded from turn-nyc-edge-vdmdp.louper.io by Hughes & Torres
Important Formulas For Algebra 2

INTRODUCTION: MASTERING THE CORE FORMULAS OF ALGEBRA 2

Algebra 2 is often seen as the bridge between foundational algebraic concepts and the more abstract world of upper-level mathematics. It is a course that challenges students to think flexibly about numbers, variables, and the relationships that govern them. Whether you are preparing for an exam, helping a child with homework, or simply brushing up on your math skills, having a solid grasp of the most important formulas for Algebra 2 can transform a daunting subject into a manageable and even enjoyable puzzle. These formulas are not just abstract rules; they are powerful tools that allow you to model real-world situations, from calculating investment growth to understanding the trajectory of a basketball. In this article, we will explore the essential formulas that every Algebra 2 student should know, broken down by topic for easy reference. By the end, you will have a clear, organized mental map of the equations that form the backbone of this critical course.

Before diving into more complex topics, it is vital to revisit the linear relationships that continue to appear throughout Algebra 2. Linear functions are the simplest type of function, and their formulas serve as the foundation for everything that follows.

The Slope-Intercept Form

The most recognizable form of a linear equation is the slope-intercept form: $y = mx + b$. Here, m represents the slope, which tells you how steep the line is, and b is the y-intercept, the point where the line crosses the y-axis. This formula is your go-to for quickly graphing a line or understanding its rate of change.

Point-Slope Form

When you know a line's slope and one point on the line, the point-slope form is incredibly useful: $y - y_1 = m(x - x_1)$. This formula allows you to write the equation of a line without needing the y-intercept directly, making it ideal for problems where you are given two points or a slope and a point.

Standard Form

Another common representation is the standard form: $Ax + By = C$, where A, B, and C are integers, and A is non-negative. While not as immediately intuitive for graphing, the standard form is excellent for solving systems of equations using elimination methods.

The Distance and Midpoint Formulas

Two geometric formulas that frequently appear in Algebra 2 coordinate geometry problems are the distance formula and the midpoint formula.

- **Distance Formula:** $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$. This is derived from the Pythagorean theorem and gives the straight-line distance between any two points in the coordinate plane.
- **Midpoint Formula:** $M = ((x_1 + x_2)/2, (y_1 + y_2)/2)$. This calculates the point exactly halfway between two given points.

QUADRATIC FUNCTIONS AND EQUATIONS

Quadratics are the heart of Algebra 2. Mastering the formulas below will allow you to solve any quadratic equation and understand the behavior of parabolic graphs.

The Quadratic Formula

Perhaps the most famous formula in all of algebra, the quadratic formula solves any quadratic equation of the form $ax^2 + bx + c = 0$:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

The expression under the square root, $b^2 - 4ac$, is called the discriminant. It tells you how many real solutions exist: if the discriminant is positive, there are two distinct real roots; if zero, one real repeated root; and if negative, two complex (non-real) roots.

Vertex Form of a Quadratic

To find the vertex of a parabola quickly, use vertex form: $y = a(x - h)^2 + k$. The vertex is at the point (h, k). This form is essential for graphing transformations of the basic parabola $y = x^2$.

Standard Form Vertex Formula

If you have a quadratic in standard form, you can find the x-coordinate of the vertex using $x = -b / (2a)$. Substituting this value back into the equation gives you the y-coordinate of the vertex.

Factoring Patterns

Several special factoring formulas appear repeatedly in Algebra 2:

- **Difference of Squares:** $a^2 - b^2 = (a - b)(a + b)$
- **Perfect Square Trinomial:** $a^2 + 2ab + b^2 = (a + b)^2$ and $a^2 - 2ab + b^2 = (a - b)^2$
- **Sum and Difference of Cubes:** $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$ and $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

POLYNOMIAL FUNCTIONS AND OPERATIONS

Beyond quadratics, Algebra 2 introduces higher-degree polynomials. Key formulas help with dividing, factoring, and analyzing these functions.

Remainder Theorem and Factor Theorem

When a polynomial $f(x)$ is divided by $(x - c)$, the remainder is $f(c)$. This is the **Remainder Theorem**. The **Factor Theorem** states that $(x - c)$ is a factor of $f(x)$ if and only if $f(c) = 0$. These theorems are fundamental for finding roots and simplifying polynomials.

Rational Root Theorem

To find possible rational roots of a polynomial with integer coefficients, the rational root theorem is extremely helpful: possible rational roots are of the form $\pm$ (factors of the constant term) / (factors of the leading coefficient). This narrows down the candidates for synthetic division.

Synthetic Division

While not a formula per se, the process of synthetic division is a quick method to divide a polynomial by a binomial of the form $(x - c)$. Knowing how to apply it efficiently is just as important as any formula.

General Form of a Polynomial

A polynomial function is often written as $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$ where $a_n \neq 0$. The degree n determines the maximum number of real roots and the general shape of the graph.

EXPONENTIAL AND LOGARITHMIC FUNCTIONS

Exponential growth and decay, along with their inverse functions—logarithms—are central to Algebra 2. These formulas

are used in finance, biology, and physics.

Exponential Growth and Decay

The basic exponential function is $y = ab^x$, where a is the initial value, b is the growth factor ($b > 1$ for growth, $0 < b < 1$ for decay), and x is the exponent. For continuous growth, use $A = Pe^{rt}$, where P is the principal, r is the rate, and t is time.

Logarithm Definition

A logarithm answers the question: to what exponent must a base be raised to get a certain number? The definition: $\log_a(x) = y$ means $a^y = x$. This equivalence is crucial for solving exponential equations.

Properties of Logarithms

Three key properties allow you to rewrite logarithmic expressions:

1. **Product Rule:** $\log_a(MN) = \log_a(M) + \log_a(N)$
2. **Quotient Rule:** $\log_a(M/N) = \log_a(M) - \log_a(N)$
3. **Power Rule:** $\log_a(M^p) = p \cdot \log_a(M)$

Change of Base Formula

When you need to evaluate a logarithm in a different base (e.g., on a calculator that only has log base 10 or natural log), use: $\log_a(x) = \log_x(x) / \log_x(a)$, where the new base can be any convenient base.

SEQUENCES AND SERIES

Algebra 2 often includes the study of arithmetic and geometric sequences and their sums. These formulas are essential for pattern recognition and real-world applications like loan payments or population growth.

Arithmetic Sequences

An arithmetic sequence has a constant difference d between terms. The n th term is given by: $a_n = a_1 + (n - 1)d$. The sum of the first n terms is: $S_n = n/2 \times (a_1 + a_n)$ or equivalently $S_n = n/2 \times [2a_1 + (n - 1)d]$.

Geometric Sequences

A geometric sequence has a constant ratio r between terms. The n th term is: $a_n = a_1 \cdot r^{n-1}$. The sum of the first n terms is: $S_n = a_1(1 - r^n) / (1 - r)$ for $r \neq 1$. For an infinite geometric series where $|r| < 1$, the sum is: $S = a_1 / (1 - r)$.

SYSTEMS OF EQUATIONS AND MATRICES

Solving multiple equations simultaneously is a key skill, and Algebra 2 often introduces matrices as a powerful tool.

Solving Linear Systems

Three main methods: substitution, elimination, and graphing. The formulas for elimination often rely on rearranging equations to the standard form $Ax + By = C$. For three variables, the method extends logically.

Cramer's Rule

For a system of linear equations represented in matrix form, Cramer's Rule uses determinants to find each variable. For a 2×2 system: $x = \det(A_x) / \det(A)$ and $y = \det(A_y) / \det(A)$, where A is the coefficient matrix and A_x , A_y are matrices with the constant column substituted.

Determinant of a 2×2 Matrix

The determinant of matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is $ad - bc$. For a 3×3 matrix, use the more involved formula using minors and cofactors.

PROBABILITY AND STATISTICS

Algebra 2 frequently includes an introduction to probability and combinatorics. These formulas quantify uncertainty and counting.

Permutations

The number of ways to arrange r items from a set of n distinct items, where order matters, is: ${}_nP_r = n! / (n - r)!$.

Combinations

The number of ways to choose r items from n , where order does not matter, is: ${}_nC_r = n! / [r!(n - r)!]$.

Probability of an Event

The probability of an event A is: $P(A) = (\text{number of favorable outcomes}) / (\text{total number of outcomes})$. For independent events, $P(A \text{ and } B) = P(A) \times P(B)$. For mutually exclusive events, $P(A \text{ or } B) = P(A) + P(B)$.

Expected Value

For a random variable X with possible values x_i and probabilities p_i , the expected value is: $E(X) = \sum (x_i \cdot p_i)$.

CONIC SECTIONS

Conic sections—parabolas, circles, ellipses, and hyperbolas—are

explored in Algebra 2. Each has a standard equation formula that defines its shape and orientation.

Circle

The equation of a circle with center (h, k) and radius r is