
Bones Of The Hand And Wrist Anatomy

*Bones Of The Hand And
Wrist Anatomy*

*Downloaded from [turn-nyc-
edge-vdmdp.louper.io](https://turn-nyc-edge-vdmdp.louper.io) by
Chance & Jeremiah*

INTRODUCTION TO THE COMPLEX ARCHITECTURE OF THE HAND AND WRIST

The human hand is one of the most intricate and versatile structures in the body, capable of everything from delicate micro-movements to powerful gripping actions. At the core of this remarkable functionality lies the **bones**

of the hand and wrist anatomy, a sophisticated arrangement of 27 individual bones that work together to provide both stability and extraordinary range of motion. Understanding the **bones of the hand and wrist anatomy** is essential not only for medical professionals but also for anyone interested in how the body achieves such precise and powerful manual dexterity.

The wrist and hand represent the distal

THE CARPAL BONES: THE connecting the forearm to the fingertips. This anatomical region is divided into three main segments: the carpus (wrist), the metacarpus (palm), and the phalanges (fingers). Each segment plays a distinct role in the overall function of the hand, and the **bones of the hand and wrist anatomy** are arranged in a way that maximizes both strength and flexibility. The wrist alone contains eight small carpal bones arranged in two rows, while the palm comprises five metacarpal bones that radiate outward to support the fingers. The fingers themselves are made up of fourteen phalanges, with each finger containing three phalanges except the thumb, which has two.

FOUNDATION OF WRIST MOTION

When examining the **bones of the hand and wrist anatomy**, the carpal bones serve as the critical link between the forearm and the hand. These eight small, irregularly shaped bones are arranged in two transverse rows, each containing four bones. The proximal row sits closer to the forearm, while the distal row connects to the metacarpal bones of the palm. Understanding these bones and their interrelationships is fundamental to grasping how the wrist achieves its impressive range of motion while maintaining structural integrity.

The Proximal Row of Carpal Bones

The proximal row consists of four bones that articulate with the radius and ulna of the forearm. From the lateral (thumb) side to the medial (pinky) side, these bones are the scaphoid, lunate, triquetrum, and pisiform. Each bone in this row has unique characteristics and functions within the **bones of the hand and wrist anatomy**.

The scaphoid bone is the largest bone in the proximal row and is shaped somewhat like a boat, which is reflected

in its name derived from the Greek word for boat. The scaphoid sits at the base of the thumb and is the most commonly fractured carpal bone due to its position and the forces transmitted through it during falls onto an outstretched hand. Its blood supply is particularly vulnerable, entering at the distal end, which means fractures can lead to avascular necrosis if not properly treated.

The lunate bone is crescent-shaped, as its name suggests, and sits between the scaphoid and triquetrum. It articulates with the radius and is central to wrist motion. The lunate is important in the **bones of the hand and wrist anatomy** because it acts as a keystone in the proximal row, and its dislocation can lead to significant wrist dysfunction.

The triquetrum bone is pyramidal in shape and sits on the medial side of the proximal row. It articulates with the lunate, the pisiform, and the hamate bone of the distal row. The triquetrum is the second most commonly fractured carpal bone and plays a role in the complex movements of wrist rotation.

The pisiform bone is a small, pea-shaped bone that sits on the palmar surface of the triquetrum. It is a sesamoid bone, meaning it develops within a tendon, specifically the flexor carpi ulnaris tendon. The pisiform acts as a pulley for the tendon and is an important landmark in the **bones of the hand and wrist anatomy** for understanding the attachment points of wrist flexors.

The Distal Row of Carpal Bones

The distal row of carpal bones connects the proximal row to the metacarpal bones of the palm. From lateral to medial, these bones are the trapezium, trapezoid, capitate, and hamate. Each bone in this row has distinct articulations and functions that contribute to the overall mechanics of the hand.

The trapezium bone is distinguished by its saddle-shaped articulation with the first metacarpal bone of the thumb. This unique saddle joint is what allows the thumb its remarkable range of motion, including opposition, which is essential for grasping and pinching. The trapezium

is a key component in the **bones of the hand and wrist anatomy** because it directly supports thumb function, and arthritis at this joint is a common source of hand pain, particularly in older adults.

The trapezoid bone is the smallest bone in the distal row and is wedge-shaped, fitting between the trapezium and the capitate. It articulates with the second metacarpal bone and contributes to the stability of the index finger ray. While less commonly discussed in the **bones of the hand and wrist anatomy**, the trapezoid plays an essential role in transmitting forces through the hand.

The capitate bone is the largest carpal bone and sits at the center of the wrist, acting as a keystone for the entire carpal arch. It articulates with the third

metacarpal bone and is involved in the majority of wrist movements. The capitate head articulates with the lunate in the proximal row, and this joint is critical for wrist flexion and extension. Understanding the capitate is essential when studying the **bones of the hand and wrist anatomy** because of its central role in wrist mechanics.

The hamate bone is wedge-shaped and sits on the medial side of the distal row. It has a distinctive hook-like projection called the hook of the hamate, which serves as an attachment point for ligaments and tendons. The hamate articulates with the fourth and fifth metacarpal bones, supporting the ring and little fingers. The hook of the hamate is a clinically important structure in the **bones of the hand and wrist**

anatomy because it can be fractured by repetitive impact, such as from golf clubs or baseball bats.

THE METACARPAL BONES: THE FRAMEWORK OF THE PALM

Moving from the wrist into the hand itself, the five metacarpal bones form the skeleton of the palm. Each metacarpal bone consists of a base, a shaft, and a head, and they are numbered from one to five, starting with the thumb. The metacarpals are long bones, and their structure reflects their role in transmitting force from the wrist to the fingers while also providing the rigid framework for the palm. In the **bones of the hand and wrist anatomy**, the metacarpals are the bridge between the carpus and the

phalanges.

The first metacarpal bone, associated with the thumb, is shorter and thicker than the others and has a unique saddle-shaped base that articulates with the trapezium. This specialized articulation allows the thumb to move in multiple planes, including opposition, which is a hallmark of human hand function. The second and third metacarpals are the most stable, articulating firmly with the trapezoid and capitate respectively, and they form the fixed central column of the hand. The fourth and fifth metacarpals have increasing mobility, which contributes to the cupping motion of the palm during grasping.

The heads of the metacarpal bones form the knuckles when the fist is clenched. These heads articulate with the proximal

phalanges of the fingers at the metacarpophalangeal joints, which are critical for finger flexion and extension. In the **bones of the hand and wrist anatomy**, the metacarpal heads are important landmarks for both function and injury, as fractures of the metacarpal neck and head are common in hand trauma.

THE PHALANGES: THE BONES OF THE FINGERS

The phalanges are the small bones that make up the fingers, and there are fourteen in total within the **bones of the hand and wrist anatomy**. Each finger, except the thumb, contains three phalanges: the proximal phalanx, the middle phalanx, and the distal phalanx. The thumb contains only a proximal and

distal phalanx, lacking a middle phalanx. The phalanges are long bones, each with a base, a shaft, and a head, and they decrease in size from proximal to distal.

The proximal phalanges articulate with the metacarpal heads at the metacarpophalangeal joints, forming the knuckles. The middle phalanges articulate with the proximal phalanges at the proximal interphalangeal joints, and the distal phalanges articulate with the middle phalanges at the distal interphalangeal joints. The distal phalanges are flattened and expanded at their tips to support the fingernails and the highly sensitive fingertip pulp. In the **bones of the hand and wrist anatomy**, the distal phalanges are also the sites of attachment for the flexor and extensor tendons, making them crucial

for fine motor control.

The thumb phalanges are particularly important in the **bones of the hand and wrist anatomy** because they contribute to the thumb's unique functional capabilities. The proximal phalanx of the thumb articulates with the first metacarpal at a condylar joint that allows flexion and extension, while the distal phalanx provides the broad surface needed for opposition and pinch strength.

JOINTS AND ARTICULATIONS: HOW THE BONES WORK TOGETHER

The **bones of the hand and wrist anatomy** do not exist in isolation; they are connected by a complex network of

joints that allow for coordinated movement. The wrist joint, properly called the radiocarpal joint, is formed by the articulation of the distal radius with the scaphoid, lunate, and triquetrum of the proximal carpal row. This joint allows for flexion, extension, radial deviation, and ulnar deviation of the wrist. The midcarpal joint between the proximal and distal carpal rows adds additional motion, particularly in the coronal plane.

The carpometacarpal joints connect the distal carpal row to the bases of the metacarpal bones. The first carpometacarpal joint, involving the trapezium and the first metacarpal, is a highly mobile saddle joint that is unique to the thumb. The second and third carpometacarpal joints are relatively rigid, providing stability for precision

LIGAMENTS AND SOFT TISSUE fifth carpometacarpal joints have moderate mobility, allowing the palm to cup during grasping.

The metacarpophalangeal joints are condyloid joints that allow flexion, extension, abduction, and adduction of the fingers. These joints are critical for gripping and manipulating objects. The interphalangeal joints are hinge joints that permit only flexion and extension, and they are essential for fine motor control during activities such as writing, typing, and pinching. Understanding these articulations is central to appreciating how the **bones of the hand and wrist anatomy** function as a cohesive unit.

CONNECTIONS

While the **bones of the hand and wrist anatomy** provide the rigid framework, ligaments and soft tissues are what hold the bones together and guide their movement. The wrist is stabilized by a complex network of intrinsic and extrinsic ligaments that connect the carpal bones to each other and to the radius and ulna. The scapholunate ligament, for example, connects the scaphoid and lunate bones and is a common site of injury in wrist sprains. The triangular fibrocartilage complex stabilizes the distal radioulnar joint and the ulnar side of the wrist.

In the hand, the volar plates are thick fibrocartilaginous structures on the

palmar side of the metacarpophalangeal and interphalangeal joints that prevent hyperextension. The collateral ligaments on the sides of these joints provide lateral stability. The transverse metacarpal ligament connects the heads of the metacarpal bones and helps maintain the transverse arch of the palm. All of these soft tissue structures work in concert with the **bones of the hand and wrist anatomy** to produce smooth, coordinated, and powerful hand function.

COMMON INJURIES AND CLINICAL RELEVANCE

Understanding the **bones of the hand and wrist anatomy** is essential for diagnosing and treating the many injuries that affect this region. Fractures of the carpal bones are common, with the scaphoid being the most frequently fractured. Scaphoid fractures often result from a fall onto an outstretched hand and can be difficult to diagnose on initial X-rays.